

IN PEOPLE WITH TYPE 1 DIABETES

DIABETIC KETOACIDOSIS (DKA)

IS DANGEROUS

Once the DKA process begins, things can progress quickly

CONFUSION



COMA




DEATH



YOU CAN TAKE ACTION TO PREVENT ITS PROGRESSION: MEASURE YOUR KETONES


* Always have ketone testing supplies on hand and, use if you're . . .

*Refer to reference page for ketones & DKA definition



Feeling sick or unwell


AND



Have very high sugars


**Refer to reference page for exceptions

Ketones negative or below 0.6? Ketones are not the problem. Recheck in 2h*



*Refer to reference page for next steps

Ketones positive or above 0.6? The process has started. You are in danger.



TAKE ACTION


IMMEDIATE ACTION

- Feeling worse
- Ketones rising
- Ketones are above 3.0
- If you are not sure what to do...

IF STILL NO IMPROVEMENT

- Stay hydrated
- Take 1.5X usual correction bolus of fresh rapid insulin by syringe or change pump site and use fresh insulin*
- Check ketones every 2 hours; repeat until ketones are below 0.6
- Never stop insulin (even if unable to eat)

*unsure of how to calculate your correction bolus? Go to the reference page!



GO TO THE HOSPITAL

You may not like the hospital, but it can save your life!

DKA REFERENCE PAGE

Be prepared: Always keep ketone strips on hand!

Keeping unexpired ketone strips on hand is vital so you can check your levels any time you're feeling unwell and have sugars above 14.0 mmol/L. Blood ketone is preferred over urine ketone, but either is a reasonable option. If you don't have any on hand, speak with your doctor about getting a prescription for ketone strips. Alternatively, you can go to a pharmacy that carries ketone strips and buy them out-of-pocket.

What are ketones?

When your body doesn't have enough insulin, your cells can't use sugar as a source of energy. As a result, your body breaks down fat to create energy instead. Ketones are an acid that your body produces as a byproduct when it breaks down fat.

What is DKA?

DKA, Diabetic Ketoacidosis, is a life threatening condition that occurs when you have too much ketones. In DKA, the excess ketones cause the blood to be more acidic than the body can handle. It usually happens when your sugars are very high.

Symptoms of DKA can be thought of as those that are due to high blood sugars and those that are due to blood acidity.

- **Hyperglycemia:** excessive thirst, dry mouth, frequent urination, blurry vision
- **Acidity:** nausea/vomiting, abdominal pain, weakness, confusion, fruity-smelling breath, shortness of breath or chest pain.

What puts you at risk of developing DKA?

Ketones develop when your body doesn't have enough insulin. Sometimes this can happen because of missed doses or failed insulin therapy (pump occlusions, bad insulin). Other times, your insulin requirements might be higher than usual because of a health stressor like infection, surgery, or other significant health events like a heart attack. Rarely, people with T1D might be put on some drugs like SGLTi's off-label. These medications increase your risk of DKA significantly.

How to calculate a correction bolus when ketones are above 0.6mmol/L:

If you usually estimate how much insulin you'd take to correct your sugar, take that guess and multiply it by 1.5 to calculate your correction for ketones. If you use an insulin pump, you should already have a correction factor, know as an insulin sensitivity factor (ISF); have the pump calculate your correction dose and multiply that by 1.5. Pumps sometimes fail, so make sure to give your dose by injection using fresh insulin. Change the pump site, tubing, and reservoir and use fresh insulin. If using Hybrid Closed Loop, you may need to temporarily come out of the Automatic Algorithm Mode. If you're unsure of how to treat a high blood sugar, it's a good idea to get in touch with your diabetes educator or go to the hospital.

DKA averted! Now what?

In the near future, make sure to follow up with your diabetes care team and let them know about this event. Things may be fine, or you may need a higher daily dose of insulin for the future.

**Exceptions related to ketone measurements:

Although not intended for use in T1D (off-label use), Sodium-Glucose Linked Transporter inhibitors (SGLTi) are associated with risk of "euglycemic DKA". In this situation, DKA can start even with normal blood sugars. If using SGLTi, ketones should be checked if feeling unwell, regardless of blood sugar levels. If ketones are positive, people should **Stop** SGLTi, take additional **Insulin**, ingest **Carbohydrates** and **Hydrate** ('StlCH' Protocol). Pregnancy, extreme exercise, nausea/vomiting/diarrhea (especially in children), or starvation/excess alcohol are other conditions associated with more normal blood sugar levels despite ketones.

Assessing risk: Checking some well day ketones rather than only on sick days:

Consider using your blood ketone strips on well days before they expire. We have found that if you check a weekly, well day morning ketone level over a month (4 tests) that if any result is 0.8mmol/l or higher that your risk for DKA may be high over the next months. Check with your diabetes team about possibly increasing your basal insulin doses, adjusting your insulin regimen, and make sure that you have a good sick day plan.

Please Note: Our study first learned the barriers in DKA prevention with 9 people living with type 1 diabetes (PLWT1D), 12 healthcare professionals, and 1 caregiver. Then, this core DKA prevention infographic was co-created between our research team, 20 PLWT1D, 17 healthcare professionals, 1 caregiver, an implementation scientist and media expert over a series of iterative design refinement steps. The infographic has also undergone an evaluation phase with 20 additional PLWT1D.

Verhoeff NJ, et al. Barriers to diabetic ketoacidosis prevention in adults with type 1 diabetes: Implications for education and implementation of new monitoring technologies. *Can J Diabetes*. 2026; <https://doi.org/10.1016/j.jcjd.2025.11.002>

Waniss MR, et al. A Patient-Clinician Co-Designed Infographic for Diabetic Ketoacidosis Education in Type 1 Diabetes . In revision. *Can J Diabetes*. 2026;in press

Bapat P, et al. Capillary Ketone Level and Future Ketoacidosis Risk in Patients With Type 1 Diabetes Using Sodium-Glucose Cotransporter Inhibitors. *Diabetes Care*. 2025; <https://doi.org/10.2337/dc25-0125>